

LA-UR-19-24567

Approved for public release; distribution is unlimited.

Title: Rapid Detection of Bacteremia in Human Blood

Author(s): Smith, Joshua E.

Intended for: Science Highlight

Issued: 2019-05-16

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Rapid Detection of Bacteremia in Human Blood

Bacteremia is a leading cause of death for children in sub-Saharan Africa where childhood mortality rates are the highest in the world, and its early diagnosis and treatment can save many lives. However, diagnosing bacteremia is challenging, especially when there is co-infection with diseases such as malaria and HIV. The good news is that research from Los Alamos published in the April 19th edition of Nature Scientific Reports indicates that there may be a way to greatly speed up diagnosis. In the paper, the scientists report on a method for the rapid and accurate detection of bacteremia using a technique that works by mimicking the way the human immune system recognizes pathogens.

Bacteremia is the presence of bacteria in the bloodstream and it can be either asymptomatic or result in a life-threatening infection that includes sepsis. Sepsis, which is when the body's response to infection causes damage to its own tissues and organs, is the most common cause of global childhood mortality. Microbial culture, which requires bacterial growth in a petri dish, is the current clinical standard for sepsis diagnosis and it is both slow, insensitive, and not practical for field diagnostics. Because of these limitations there is a pressing need for a rapid and accurate method that can direct therapeutic intervention at the point of need. The work presented by the Los Alamos authors is the first step towards developing such a capability and is inspired by the ability of the human immune system to sensitively detect bacteremia in blood very early in the course of infection by its reaction to pathogen biomarkers.

The researchers developed and clinically validated a novel method for the direct detection of the same pathogen biomarkers that the immune system uses to recognize bacteremia. Pathogen biomarkers are present for both Gram-negative bacteria and Gram-positive bacteria. The Gram test is a method to broadly categorize bacteria in two categories and it is important to be able to detect bacteria in both. These biomarkers interact with host lipoprotein carriers in blood, which allows for their direct detection using two tailored assays developed at Los Alamos called membrane insertion and lipoprotein capture (see figure 1). The detection occurs in a waveguide-based optical biosensor platform that was also developed at Los Alamos. The team also demonstrated the application of these methods for the detection of *Salmonella* Typhimurium bacteremia and *Staphylococcal* bacteremia. Taken together, these results demonstrate the significance of biochemistry in both understanding of host-pathogen biology, and the development of assay methodology, as well as demonstrating a method for the rapid, sensitive and accurate diagnosis of bacteremia at the point of need.

Ref: "Direct detection of bacteremia by exploiting host-pathogen interactions of lipoteichoic acid and lipopolysaccharide," *Scientific Reports* 9, Article number: 6203 (2019)
<https://www.nature.com/articles/s41598-019-42502-5>

Technical contact: Harshini Mukundan




Figure 1. Overview of bacterial pathogen associated molecular patterns (PAMP) detection strategies. (a) Schematic representation of amphiphilic bacterial biomarkers in association with biochemically similar molecules either by forming micelles or interacting with host lipoprotein carriers. Black arrows indicate physiological associations, while orange arrows indicate experimental processes. In the absence of sequestration by a host lipoprotein carrier, bacterial PAMPs can be detected by (b) membrane insertion, which requires only one antibody. When associated with a host lipoprotein carrier, detection can be performed using (c) lipoprotein capture, which requires two antibodies, as well as prior knowledge of PAMP-lipoprotein carrier associations. Dimensions of the various layers are included in order to present the biophysical dimensions of assay performance within the evanescent field of the waveguide. Graphic representations are not drawn to scale.




Figure 2. Aaron Anderson of the Physical Chemistry and Applied Spectroscopy group adjusts a field-portable version of the optical biosensor.